

Review

Immune Reconstitution after Haploidentical Hematopoietic Stem Cell Transplantation

Ying-Jun Chang¹, Xiao-Yu Zhao¹, Xiao-Jun Huang^{1,2,*}¹ Peking University People's Hospital and Peking University Institute of Hematology, Beijing Key Laboratory of Hematopoietic Stem Cell Transplantation, Beijing, China² Peking-Tsinghua Center for Life Sciences, Beijing, China

Article history:

Received 7 August 2013

Accepted 29 November 2013

Key Words:

Haploidentical transplantation

Immune reconstitution

Natural killer cell

T cell

A B S T R A C T

Haploidentical hematopoietic stem cell transplantation (HSCT) offers the benefits of rapid and nearly universal donor availability and has been accepted worldwide as an alternative treatment for patients with hematologic malignancies who do not have a completely HLA-matched sibling or who require urgent transplantation. Unfortunately, serious infections and leukemia relapse resulting from slow immune reconstitution remain the 2 most frequent causes of mortality in patients undergoing haploidentical HSCT, particularly in those receiving extensively T cell–depleted megadose CD34⁺ allografts. This review summarizes advances in immune recovery after haploidentical HSCT, focusing on the immune subsets likely to have the greatest impact on clinical outcomes. The progress made in accelerating immune reconstitution using different strategies after haploidentical HSCT is also discussed. It is our belief that a predictive immune subset–guided strategy to improve immune recovery might represent a future clinical direction.

© 2014 American Society for Blood and Marrow Transplantation.

INTRODUCTION

Haploidentical hematopoietic stem cell (HSC) transplantation (HSCT) is available for nearly all patients and has no search or acquisition costs [1–9]. Over the past decade, many haploidentical transplantation protocols, including T cell–replete and T cell–depleted (TCD) haplotype HSCT, depending on whether or not allografts have been engineered in vitro, have demonstrated promising clinical outcomes [4,8,9]. Unfortunately, serious infections and disease relapse resulting from delayed immune reconstitution remain the 2 most frequent causes of mortality after allogeneic HSCT, particularly in patients who received extensively TCD CD34⁺ cell megadose allografts [10–15]. Advances in the understanding of immune recovery profiles in haploidentical recipients [8,16–47], along with new methods of modifying donor T cells [48–50] and natural killer (NK) cells [51], have made it possible to establish new strategies to improve post-transplantation immunologic reconstitution [9,52–58].

In this review, we summarize advances in immune recovery after haploidentical HSCT, focusing on the recovered immune subsets likely to have the greatest impact on clinical outcomes [18,23,33,42,45,59–63]. We compare the differences in immune reconstitution between haploidentical transplantation and other transplantation strategies, including HLA-identical sibling donor transplantation, HLA-matched unrelated donor transplantation, and umbilical cord blood transplantation. In addition, we discuss recent advances in the enhancement of immune reconstitution after haploidentical HSCT [64–70].

KINETICS OF IMMUNE RECOVERY AFTER HAPLOIDENTICAL HSCT

Different immune cell subgroups recover at different rates after haploidentical HSCT (Table 1). The conditioning regimen is followed by a “neutropenic” phase that lasts until neutrophils reconstitute, at a median of approximately 11 to 12 days after TCD HSCT with high doses of CD34⁺ cells [8,38], approximately 15 days after unmanipulated haploidentical steady-state bone marrow (BM) allografts, approximately 21 days after granulocyte colony-stimulating factor (G-CSF)-primed BM allografts, and approximately 13 days after G-CSF-stimulated blood and BM allografts [2,7,71]. The recovery of neutrophil function (eg, chemotaxis, phagocytosis, superoxide production, killing of bacteria) in haploidentical settings remains poorly understood, however.

The rapid recovery of NK cells after haploidentical transplantation is based on an expansion of the cytokine-producing CD56^{bright} NK cell subsets in both T cell–replete and TCD settings [16,17,24,41]. Although the absolute number of overall NK cells usually recovers to the donor's level by day 30 post-HSCT, recovery may be delayed by the development of graft-versus-host disease (GVHD) [8,24]. The function of NK cell recovery is regulated by the balance of activating and inhibitory signals transmitted by different cell surface receptors. Thus, the expression of activating NK receptors (NKR), such as NKP46, NKP44, NKP30, and NKG2D, as well as inhibiting NKRs, such as CD158a, CD158b, CD158e, and NKG2A, is essential for NK cell activation.

After haploidentical HSCT, the overall expression of activating NKRs and inhibitory NKRs is reduced, whereas CD94/NKG2A expression is increased. NKG2A recovery is inversely correlated with CD158 recovery in the year after HSCT. This altered phenotype includes more CD56^{bright} cells, fewer CD56^{dim} NK cells, and altered CD94/NKG2A expression. Activation or inhibition of NKR expression during early reconstitution is associated with lower levels of in vitro NK

Financial disclosure: See Acknowledgments on page 447.

* Correspondence and reprint requests: Xiao-Jun Huang, MD, Professor, Peking University Institute of Hematology, Peking University People's Hospital, No. 11 Xizhimen South St, Beijing 100044, China.

E-mail address: xjhrm@medmail.com.cn (X.-J. Huang).

1083-8791/\$ – see front matter © 2014 American Society for Blood and Marrow Transplantation.

<http://dx.doi.org/10.1016/j.bbmt.2013.11.028>

cytotoxicity after haploidentical HSCT. The reconstitution of killer cell immunoglobulin-like receptors (KIRs) is influenced by many factors, including the conditioning regimen, level of T cell depletion, and the use of immune suppression after transplantation, as recently reviewed by Zhao et al. [72]. Monocyte engraftment is rapid, with normal values on day +15 after unmanipulated haploidentical blood and BM transplantation [25]. In HLA-identical HSCT settings, the absolute monocyte counts at day +30 (>300 cells/ μ L) are strongly associated with improved survival [73,74], whereas no association between monocyte recovery and outcome after haploidentical HSCT has been reported.

Dendritic cells (DCs) are highly important antigen-presenting cells with central roles in initiating and modulating immune responses. Human peripheral blood DCs are divided into 2 major subsets: myeloid DCs (mDCs) and plasmacytoid DCs (pDCs). mDCs are further subdivided into 2 subsets: T helper cell (Th) 1-promoting mDCs (mDC1s) and Th2-promoting mDCs (mDC2s) [75]. Our preliminary data showed extremely low mDC1, mDC2, and pDC counts at 1–3 months post-HSCT, which recovered to normal values by 1 year after unmanipulated haploidentical blood and BM transplantation [25]. A gradual increase in the DC population after transplantation was also reported in pediatric patients with acute leukemia who received CD3/CD19-depleted grafts from haploidentical donors [41]. The early delayed immune reconstitution of DCs may contribute in part to an increased rate of postengraftment bacterial and fungal infections [76], although little is known about the recovery of DC function after haploidentical transplantation [25,41].

Invariant NK T (iNKT) cells are a specialized subset of T cells that use their T cell receptors (TCRs) to recognize self and foreign lipids presented by CD1d as cognate antigens. These cells can have protective or harmful roles in many pathological states, including microbial infection, autoimmune disease, allergic disease, and cancer [77]. In pediatric patients, de Lalla et al. [33] observed that iNKT recovered slowly after CD34-selected haploidentical HSCT and reached normal reference values by 18 months post-transplantation. Their data also suggest that the frequency of iNKT cells is significantly correlated with the remission state after haploidentical HSCT. In HLA-identical transplantation settings, Chaidos et al. [78] reported that CD4(-) iNKT cell dose was the sole graft parameter predictive of clinically significant acute GVHD. This interesting finding awaits confirmation in haploidentical HSCT settings.

Detailed analyses of adaptive immune reconstitution have been performed by researchers at various transplantation centers [16,17,22,25–27,29,32,33,38,41,45,79–81]. After CD34-selected haploidentical HSCT, mean (\pm standard deviation) CD4⁺ cell counts ranged from 100 ± 40 /L to 200 ± 20 /L at 10 months post-transplantation, and rose thereafter. The mean CD8⁺ cell count reached 230 ± 80 /L on day +60, followed by a steady rise to 570 ± 125 /L by day +300. The mean CD16⁺ NK cell count reached 400/L stably by day +30 [38]. The recovered NK cells may enhance the graft-versus-leukemia effect [38–82]. A German group used negative selection of CD3/CD19 cells to retain NK cells, monocytes, and DCs in haploidentical allografts and found delayed T cell reconstitution, with a median of 369 CD3⁺ cells/ μ L, 177 CD4⁺ cells/ μ L, and 193 CD8⁺ cells/ μ L on day +400 [8,83]. In our unmanipulated haploidentical blood and BM transplantation protocol, a median absolute number of 277 CD4⁺ T cells/ μ L and 884 CD8⁺ T cells/ μ L were recovered at 1 year post-transplantation [25]. Immune recovery

after unmanipulated haploidentical HSCT appears to be more rapid than CD34-selected haploidentical transplantation; however, the 2 transplantation methods are not strictly comparable [17,25,38,39].

The expansion of memory T cells (especially CD4⁺ memory T cells, which reconstitute later than CD8⁺ memory T cells and rely more on thymic production of naive T cells) results in a significant inversion of the CD4/CD8 ratio up to 1 year after transplantation [25,39,41]. In contrast to most previously reported results [25,27,36,38,39,41], no significant inversion of the CD4/CD8 ratio was observed after haploidentical HSCT using reduced-intensity conditioning and CD3/CD19-depleted grafts [17]. In the TCD haploidentical HSCT protocol [45], the main lymphocyte populations (including CD3⁺ T cells, CD4⁺ T cells, CD8⁺ T cells, and B cells) and the CD4/CD8 ratio reconstituted to age-matched control levels by 4 to 6 years post-transplantation. Similar kinetics in B cell recovery have been reported by our group and others [17,25,41,84], with absolute numbers of B cells increasing gradually and reaching normal levels more than 1 year after HSCT.

Regarding the reconstitution of T cell function, an interesting finding is that the ability of T cells to secrete IFN and IL-4 recovers to normal level by day +30 post-HSCT in patients without acute GVHD, although TCR rearrangement excision DNA circle (TREC) levels remain low for 12 to 24 months after unmanipulated haploidentical HSCT [29]. At 4 to 6 years after TCD haploidentical HSCT, recipients demonstrate higher proportions of CD31⁺ naive CD4⁺ T cells compared with donors [17,45]. The signal-joint TCR excision circle (sjTREC) levels in the recipients tend to be higher than the levels in donors, which are similar to those seen in age-matched control subjects.

The foregoing observations suggest that long-term T cell reconstitution is critically influenced by de novo T cell production, reflecting thymopoiesis as a central mechanism. A skewed and restricted distribution of TCR repertoires has been observed at less than 180 days post-transplantation [17]. Azevedo et al. [45] reported that at 4–6 years post-transplantation, patient CD4⁺ T cells and CD8⁺ T cells exhibit a diverse T cell repertoire and reconstitute a diverse regulatory T cell (Treg) pool through a mechanism involving de novo thymic production. Their findings provide strong evidence that a normal immune system can be reconstituted after haploidentical HSCT; however, the small number of patients in their study precluded from a subgroup analysis to identify the factors influencing immune reconstitution. In addition, the function of T cells in unmanipulated haploidentical HSCT remains to be investigated in depth [2,71,85–88].

More recently, Ciurea et al. [36] demonstrated that compared with patients who underwent TCD haploidentical HSCT, patients who received T cell–replete haploidentical allografts exhibit better immune reconstitution of T cell subsets, including CD4⁺ T cells, CD8⁺ T cells, naive T cells, and memory T cells, during the first 6 months after transplantation (Table 1). The authors suggested that improved immune reconstitution might contribute to the improved early outcomes observed with the use of T cell–replete grafts [36]. Thus, a prospective study to compare immune recovery between TCD and T cell–replete haploidentical approaches is warranted.

Overall, the first 90 days after haploidentical HSCT are characterized by persistent CD4⁺ and CD4⁺ naive T cells, B cell lymphopenia, and low thymic function, which render patients especially susceptible to viral and fungal infections

Table 1
Immune Reconstitution in Haploidentical Transplantation and MSDT, MUDT, and Umbilical Cord Blood Transplantation

| | Federmann et al. [17] | Chang et al. [25] | | Ciurea et al. [36] | | Pérez-Martínez et al. [41] | | Kesserwan et al. [43]* | Dominietto A et al. [93]* | | | |
|--|--------------------------|-------------------|----------------------|--------------------|----------------|----------------------------|-----------|---------------------------|---------------------------|-----------|----------------|-----------|
| | | MSDT | HBMT | TCD | T Cell Replete | HIT | MUDT | | ALT | CBIBT | HIT | MSDT |
| No. of patients | 28 | 25 | 50 | 33 | 32 | 15 | 15 | 21 | 125 | 103 | 40 | 176 |
| Age, yr, median (range) | 45 (19–65) | 38.5 (24–57) | 24.5 (7–48) | 36 (18–56) | 45 (20–63) | 9.7 ± 1.1 | 8.6 ± 1.4 | 45 (29–46) | NA | NA | NA | NA |
| Male/female sex, n | 15/13 | 17/8 | 28/22 | 16/17 | 19/13 | 14/1 | 11/4 | NA | NA | NA | NA | NA |
| Diagnosis | HM | HM | HM | HM | HM + others | AML/ALL | AML/ALL | HM | NA | NA | NA | NA |
| Myeloablative conditioning | 0 | Modified Bu/Cy | Modified Bu/Cy + ATG | 33 | 26 | 0 | 0 | Cy + TBI + Flu | NA | NA | NA | NA |
| Reduced-intensity conditioning | Flu + Mel + OKT3 + Thi | 0 | 0 | 0 | 6 | 15 | 15 | 0 | NA | NA | NA | NA |
| GVHD prophylaxis | MMF | MTX + CSA + MMF | MTX + CSA + MMF | ATG | Cy + Tac + MMF | CSA | CSA | Cy + CSA + MMF | ATG | CSA + MMF | CSA + Cy + MMF | CSA + MTX |
| Allografts | CD3/CD19-depleted G-PB | G-BM + G-PB | G-BM + G-PB | SS-BM | SS-BM | CD3/CD19-depleted G-PB | SS-BM | SS-BM | SS-BM | SS-BM | SS-BM | SS-BM |
| Immune recovery, cells/L | | | | | | | | | | | | |
| CD3 ⁺ T cells | | | | | | | | | | | | |
| Day 30 post-SCT | 8 at day +40 | 348 | 135 | NA | NA | 870 | 226 | 31 | 146 | 30 | 195 | 477 |
| Day 90 post-SCT | 205 at day +100 | 1048 | 883 | NA | NA | 983 | 652 | 413 at day +60 | 404 | 57 | 182 | 565 |
| Day 180 post-SCT | 369 at day +400 | 1037 | 1163 | NA | NA | NA | NA | 691 | 470 | 196 | 499 | 700 |
| CD3 ⁺ CD4 ⁺ T cells | | | | | | | | | | | | |
| Day 30 post-SCT | 5 at day +40 | 141 | 21 | 1 | 26 | 141 | 138 | 11 | 36 | 7 | 45 | 160 |
| Day 90 post-SCT | 70 at day +100 | 220 | 152 | 7 | 127 | 305 | 198 | 154 at day +60 | 86 | 36 | 127 | 170 |
| Day 180 post-SCT | 177 at day +400 | 276 | 163 | 69 | 194 | NA | NA | 193 | 111 | 106 | 211 | 198 |
| CD3 ⁺ CD8 ⁺ T cells | | | | | | | | | | | | |
| Day 30 post-SCT | 2 at day +40 | 115 | 98 | 1 | 22 | 730 | 85 | 11 | 102 | 42 | 73 | 280 |
| Day 90 post-SCT | 66 at day +100 | 645 | 672 | 17 | 181 | 678 | 447 | 193 at day +60 | 278 | 16 | 424 | 389 |
| Day 180 post-SCT | 193 at day +400 | 580 | 918 | 54 | 167 | NA | NA | 409 | 413 | 51 | 408 | 500 |
| CD3 ⁺ CD4 ⁺ naïve cells | | | | | | | | | | | | |
| Day 30 post-SCT | NA | 90 | 6 | 0.7 | 3.45 | 3.9 | 5.3 | 0.45 | NA | NA | NA | NA |
| Day 90 post-SCT | 28 at day +100 | 152 | 33 | 0.55 | 7.3 | 3.3 | 3.3 | 9 at day +60 | NA | NA | NA | NA |
| Day 180 post-SCT | 115 at day +400 | 196 | 69 | 4.8 | 6.72 | NA | NA | 24 | NA | NA | NA | NA |
| CD3 ⁺ CD4 ⁺ memory cells | | | | | | | | | | | | |
| Day 30 post-SCT | NA | 131 | 20 | 2.6 | 24 | 136 | 134 | 4 | NA | NA | NA | NA |
| Day 90 post-SCT | 79 at day +100 | 155 | 108 | 16 | 97 | 299 | 181 | 65 at day +60 | NA | NA | NA | NA |
| Day 180 post-SCT | 205 at day +400 | 261 | 171 | 111 | 179 | NA | NA | 48 | NA | NA | NA | NA |
| B cells | | | | | | | | | | | | |
| Day 30 post-SCT | NA | 13 | 13 | 5 | 0.95 | 39 | 24.6 | 0.5 | NA | NA | NA | NA |
| Day 90 post-SCT | 100 at day +150 | 33 | 23 | 55 | 64 | 147 | 43 | 119 at day +60 | NA | NA | NA | NA |
| Day 180 post-SCT | 275 at day +400 | 60 | 54 | 58 | 95 | NA | NA | 104 | NA | NA | NA | NA |
| NK cells | | | | | | | | | | | | |
| Day 30 post-SCT | 248 at day +20 | NA | NA | 225 | 84 | 291 | 254 | 59 | NA | NA | NA | NA |
| Day 90 post-SCT | NA | NA | NA | 357 | 183 | 299 | 253 | 189 at day +60 | NA | NA | NA | NA |
| Day 180 post-SCT | NA | NA | NA | 290 | 172 | NA | NA | 168 | NA | NA | NA | NA |

HBMT indicates haploidentical blood and BM transplantation; HM, hematologic malignancy; AML, acute myelogenous leukemia; ALL, acute lymphoblastic leukemia; ALT, alternative donor (1 antigen-mismatched family or unrelated donors) transplantation; CBIBT, unrelated cord blood grafted intrabone transplantation; NA, not available; Cy, cyclophosphamide; TBI, total body irradiation; Flu, fludarabine; Bu, busulfan; ATG, anti-human thymocyte immunoglobulin; Mel, melphalan; Thi, thiotepa; MMF, mycophenolate mofetil; CSA, cyclosporine A; MTX, methotrexate; Tac, tacrolimus; G-BM, G-CSF–primed BM; G-PB, G-CSF–mobilized peripheral blood stem cell grafts; SS-BM, steady-state BM.

* Data coming from meeting abstract.

Table 2
Effect of Recovered Immune Subsets on Outcomes after Haploidentical Transplantation

| Disease (No. of Patients) [Reference] | Allografts | Conditioning Regimen | Reconstituted Immune Subsets | Potential Effects on Clinical Outcomes |
|---|--|-----------------------------------|--|--|
| Hematologic malignancies (27); other diseases (6) [23] | Selected CD34 ⁺ cells | MA (33) | Thymic-dependent T cells: sTREC <0.1/150,000T TREC <0.001/150,000T CD3 ⁺ CD8 ⁺ T cells | Low thymic function is associated with leukemia relapse. |
| Hematologic malignancies (30); other diseases (2) [90] | Unmanipulated BM (13) and PBSCs (6); selected CD34 ⁺ cells (13) | MA (32) | | Univariate analysis demonstrated superior survival in patients with CD8 ⁺ CD3 ⁺ absolute counts above the 5th percentile of age-matched normal levels during the first year post- transplantation. |
| Hematologic malignancies (43) [24] | G-BM + G-PB | MA (43) | CD56 ^{bright} NK cells (9.27) and T:NK ratio at day 14 post-transplantation | The patients with more CD56 ^{bright} NK cells in the recovery stage had a higher survival rate (hazard ratio [HR], 0.406; <i>P</i> = .017) and the patients with a T/NK ratio >1.0 had a greater likelihood of getting aGVHD (HR, 3.436; <i>P</i> = .059) and chronic GVHD (HR, 3.925; <i>P</i> = .028). |
| Hematologic malignancies (206) [18] | G-BM + G-PB | MA (206) | Early recovered lymphocytes | Multivariate analysis showed that patients with higher day 30 ALC (≥300/μL) was associated with low relapse rate, low TRM, and superior survival in both adult and pediatric patients. |
| Hematologic malignancies (60) [59] | G-BM + G-PB | MA (60) | Early recovered lymphocytes | Patients with a day 30 ALC >200 cells/ μL had a markedly improved overall survival and event-free survival. The frequency of iNKT cells was significantly correlated with remission after transplantation. No CMV DNAemia was evident. |
| Hematologic malignancies (78) [60] | G-BM + G-PB | MA (78) | Early recovered lymphocytes | |
| Hematologic malignancies (21) [43] | SS-BM | MA (21) | Early recovered lymphocytes | |
| Hematologic malignancies (22) [33] | G-PB + TCD | MA (22) | iNKT | A significant correlation was seen between the number of lymphoid DC2 ⁺ cells on day +60 with patient survival. |
| Hematologic malignancies (98); nonmalignancies (33) [61] | BM (78); G-PB (42); umbilical cord blood (11) | MA (131) | CMV-specific CD8 ⁺ T cells (3 cyt ⁺ cells/μL); CMV-specific CD4 ⁺ T cells (1 cyt ⁺ cells/μL) | Low TRM and virus reactivation were noted. |
| Acute leukemia (30) [41] | G-PB with CD3/CD19 depletion | MA (30) | Lymphoid DC2 ⁺ cells | Multivariate analysis showed that a cutoff ratio value of 9% yielded the most accurate predictions of future aGVHD incidence. Treg:CD4 ⁺ T cell ratios of <9% predicted a significantly higher incidence of aGVHD compared with ratios ≥9% (<i>P</i> = .0082). |
| Hematologic malignancies (89) [91] | TCD and TCR allografts | MA (89) | CMV-specific IFN ELISPOT (1000 spots/mL) | |
| Hematologic malignancies (47) [63] | Unmanipulated PBSC (27); BM (20) | Nonmyeloablative (30); MA (17) | Treg:CD4 ⁺ T cell ratio at day 14 post-transplantation | |

PBSCs indicate peripheral blood stem cells.

and leukemia relapse [2,3,71,87,88]. These results underscore the critical importance of improving immune recovery in both TCD and T cell–replete haploidentical HSCT settings.

ASSOCIATION OF RECOVERED IMMUNE CELLS WITH TRANSPLANTATION OUTCOMES

A number of studies have evaluated the association of immune recovery with clinical outcomes after haploidentical transplantation (Table 2) [18,24,59,60]. A higher absolute lymphocyte count at day 30 post-transplantation (ALC-30; ≥300 cells/μL) is associated with lower incidences of relapse and transplantation-related mortality (TRM) and superior leukemia-free survival and overall survival in both adult and pediatric patients after unmanipulated haploidentical blood and BM transplantation [18,59,60]. Using the unmanipulated haploidentical HSCT protocol, the Peking University group demonstrated a survival advantage for patients with high early (day +14) levels of CD56^{bright} NK cells after unmanipulated haploidentical HSCT [24]. In Spain, Martin et al. [89]

confirmed this result using unmanipulated haploidentical BM grafts after RIC. Other factors, including lower numbers of recovered CD3⁺CD8⁺ T cells and lymphoid mDC2 cells, may contribute to inferior survival [41–90]. Fujioka et al. [63] reported that a Treg:CD4⁺ T cell ratio <9% is predictive of a significantly higher incidence of acute GVHD compared with a ratio ≥9% (*P* = .0082) after unmanipulated blood or BM transplantation. This finding awaits confirmation in other haploidentical HSCT protocols, such as transplantation with CD3/CD19-depleted allografts [8] and BM transplantation with high-dose post-transplantation cyclophosphamide [71].

Another study investigated multiple parameters of T cell immune reconstitution in 89 patients who underwent haploidentical HSCT. Using a receiver operating characteristic curve analysis of cytomegalovirus (CMV)-specific IFN enzyme-linked immunosorbent spot (ELISPOT) assay results at day 30 to 90, Noviello et al. [91] found that cutoff values of 1000 spots/mL allowed researchers to identify those patients

who did not reactivate the virus at later time points with high specificity (>95%). Strikingly, the 2-year TRM rate was 32% in patients with an ELISPOT finding of <1000 spots/mL and 0% in those with >1000 spots/mL ($P < .05$). This interesting finding warrants the use of a predictive biomarker in larger multicenter series to investigate whether a CMV-specific IFN- γ ELISPOT cutoff value of 1000 spots/mL is a strong surrogate biomarker for TRM. When Clave et al. [23] evaluated thymic function in a group of pediatric recipients, they found that 6 months after CD34-selected HLA-haploidentical HSCT, a TREC value below the limit of detection (<0.1 per 150,000 CD3 $^{+}$ T cells for sjTREC and <0.001 per 150,000 CD3 $^{+}$ T cells for TREC) was associated with an increased risk of relapse [23].

In summary, prospective studies with larger patient populations are needed to draw definitive conclusions regarding the impact of recovered immune cells on clinical outcomes in T cell-replete and TCD nonmyeloablative or myeloablative haploidentical settings [3,8,92]. These studies are important to the prediction of such clinical outcomes as infection rates, relapse, and TRM [37,62]. Shimoni [62] suggested that the ALC-30 has long been a simple test of immune recovery that can predict outcome after autologous HSCT as well as allogeneic and haploidentical HSCT [18,59,60]. It is easy, reproducible, and requires no special expertise. Thus, the measurement of ALC-30 may have practical significance in applying interventions to facilitate immune reconstitution as described previously for haploidentical transplants [18,42,59,60].

COMPARISON OF IMMUNE RECOVERY BETWEEN HAPLOIDENTICAL AND OTHER TRANSPLANTATION MODALITIES OR GRAFT OPTIONS

Several published prospective and retrospective studies have highlighted the differences in immune reconstitution after unmanipulated haploidentical HSCT versus HLA-matched sibling donor transplantation (MSDT), matched unrelated donor transplantation (MUDT), or umbilical cord blood transplantation (Table 1). Chang et al. [25] found that compared with HLA-matched recipients, haploidentical recipients had lower counts of T cells (particularly CD4 T cells) and DC subsets before day +90, as well as greater CD28 expression on CD4 $^{+}$ and CD8 $^{+}$ T cells, whereas T cells were equally functional in the 2 patient groups at day +30. In addition, Dominietto et al. [93] recently demonstrated early delayed immune reconstitution after haploidentical transplantation (HIT) compared with MSDT [93]. Several lines of evidence suggest an association between improved immune recovery and better transplantation outcomes [13,24,94–96]; however, in our center, transplantation outcomes regarding overall survival and leukemia-free survival were comparable after HIT and MSDT, although unmanipulated HIT was characterized by delayed early reconstitution of T cells and DCs, as well as a higher incidence of CMV antigenemia [86,87]. Several factors may account for this finding. Compensatory expansion of monocytes, NK cells, and cytotoxic T lymphocytes (CTLs) [25], especially CMV-pp65 peptide-specific CTLs with the central memory CD45RO $^{+}$ CD62L $^{+}$ cell phenotype, accompanies the recovery of CD8 $^{+}$ T cells. This cell population may proliferate and differentiate into effector memory T cells stimulated with CMV antigen and may contribute to a reduced incidence of CMV disease [31,47]. Other possible contributing factors include the preemptive management of CMV antigenemia, a superior graft-versus-leukemia effect associated with transplantation of haploidentical versus

HLA-identical sibling donor grafts for high-risk acute leukemia [97], and improvements in minimal residual disease-directed relapse prophylaxis using modified donor lymphocyte infusion [66,98–100].

In a detailed prospective analysis of immune reconstitution in MUDT and HIT with CD3/CD19-depleted grafts in 30 pediatric patients, Pérez-Martínez et al. [41] found more rapid CD8 and CD8 memory T cell recovery during the first month in HIT compared with MUDT (730 ± 208 versus 85 ± 37 cells/mL and 723 ± 274 versus 83 ± 29 cells/mL, respectively; $P < .05$). CD80 expression in the DC1-DC2-subset was higher in HIT patients than the CD80 expression in the DC1-DC2-subset in MUDT patients during the first 2 months after HSCT (respectively, 26 ± 6.98 vs 5 ± 1.4 /mL, $P = .04$ and 28 ± 12 vs 2 ± 0.8 /mL, $P = .05$); however, the reconstitution of NKT cells, NK cells, and B cells was comparable in the 2 groups [41]. Szabolcs et al. [101] reported an association between CD8 $^{+}$ lymphocytosis and effector memory phenotype and IFN secretion in children who developed de novo opportunistic infections. Brown et al. [102] presented data on recent thymic emigrants and CMV-specific immune responses demonstrating associations between higher TREC and CMV-specific immune responses and improved survival. Sauter et al. [103] found that an ATG-free regimen after double cord blood transplantation could result in more rapid CD4 $^{+}$ T lymphocyte reconstitution, and reported no infection-related deaths after day +120, suggesting a correlation between immune recovery and infection-related death. Continuing new insights into the mechanisms underlying differences in the kinetics of immune reconstitution after HIT compared with MSDT and MUDT may help guide the design of useful strategies to accelerate immune recovery in HIT settings.

STRATEGIES TO IMPROVE IMMUNE RECONSTITUTION AFTER HAPLOIDENTICAL HSCT

Many of the currently available approaches to accelerating immune recovery, including IL-7, keratinocyte growth factor, growth hormone, and adoptive transfer of immune cell subsets, have been reviewed elsewhere [12,19,37,66,104–106]. Here we discuss only recent informative updated trials and reports (Table 3); data reported between 2012 and 2013) regarding the promotion of immune reconstitution, particularly in haploidentical transplantation settings [53,65,69,107].

Cytokines

IL-7

Among the cytokines used to enhance T cell reconstitution, IL-7 certainly tops the list. In a phase 1 trial, Perales et al. [65] treated 12 patients who underwent MSDT or MUDT with escalating weekly doses of IL-7 for a 3-week period. Three patients received 10 μ g/kg, 6 patients received 20 μ g/kg, and 3 patients received 30 μ g/kg. IL-7 was well tolerated overall, with only 1 patient developing acute skin GVHD. Compared with baseline values, levels of recovered CD3 $^{+}$, CD4 $^{+}$, and CD8 $^{+}$ T cells increased by 4.3-, 6.1-, and 4.3-fold, respectively. IL-7 also enhanced functional T cell responses and TCR diversity. Unfortunately, IL-7 appeared to have no significant effect on CD4 $^{+}$ CD25 $^{+}$ FoxP3 $^{+}$ T cells, NK cells, or B cells, although the data that IL-7 significantly enhances T cell recovery are exciting [65]. Based on the foregoing findings, other approaches applied alone or in combination with IL-7 to promote the reconstitution of all immune subsets merit investigation in both HLA-matched and haploidentical transplantation protocols [2,3,87,88].

Table 3
Recent Informative* Trials and Reports Regarding Strategies for Enhancing Immune Recovery

| Strategy | Transplantation Protocol | Key Points | Reference |
|--|--------------------------|---|----------------------------|
| Approaches used in preclinical model | | | |
| IL-15 | HIT mice model | IL-15 can be used for enhancing antileukemia effect of haploidentical HSCT, which requires presence of donor-derived T cells, especially CD8 ⁺ T cells. | Sauter et al. 2013 [58] |
| Depletion of naive lymphocyte with Fas ligand | HIT mice model | The proposed approach eliminates severe GVHD in T cell –replete transplants, preserves the tumor lytic activity of mismatched donor T cells, maintains T cell reactivity to third-party antigens, and, facilitates HSC engraftment. | Askenasy et al. 2013 [49] |
| Approaches used in clinical trial or report | | | |
| IL-7 | MRDT and MUDT | Treatment 12 patients with IL-7 increases T cell recovery, induces T cell responses, and enhances TCR diversity. IL-7 have no significant effect on Treg, NK or B cells. | Perales et al. 2012 [65] |
| Escalating-dose DLI | HIT | This study suggests that DLI, using an escalating-dose regimen, can be safely given to mismatched transplant recipients with a high rate of molecular remission and a low risk of GVHD. | Innes et al. 2013 [55] |
| Modified DLI | HIT and MRDT | Eight hundred and fourteen patients were enrolled, this study suggest that risk stratification-directed modified DLI may reduce relapse and improve survival of subjects with standard-risk acute leukemia after HSCT, although the effect of DLI on immune reconstitution need to be investigated. | Yan et al. 2012 [98] |
| Preemptive immunotherapy with NK cells | HIT | In this prospective phase II study, the results suggest that NK cell activation/expansion may be required to attain clinical benefit, while careful consideration must be given to the number of T cells infused. | Stern et al. 2013 [108] |
| Adoptive transfer of suicide gene-transduced donor T cells | HIT | Thymopoiesis is reactivated after the infusion of gene-modified lymphocytes, enhancing the generation of a protective T cell compartment that is able to reduce the incidence of infectious and control of GVHD. | Vaglo et al. 2012 [65] |
| Adoptive transfer of donor-derived CMV-specific T cells | HIT, MSDT, and MUDT | Infusion of CMV-specific T cell early posttransplant reduced the need for pharmacotherapy without increased rates of CMV-related organ damage and increasing acute and chronic GVHD. | Blyth et al. 2012 [69] |
| Adoptive transfer of trivirus-directed T cells | HIT, MSDT, and MUDT | In ten patients, trivirus-directed T cells administration produced completed virological responses in 80% cases and a decrease in viral load was associated with an increase in the frequency of T cell directed against the infecting virus. No toxicities were observed. | Gerdemman et al. 2013 [63] |

MRDT indicates HLA-matched related donor transplantation; DLI, donor lymphocyte infusion.

* Published between 2012 and 2013.

IL-15

Using a clinically relevant haploidentical murine transplantation model, Sauter et al. [58] found that IL-15 administration significantly increased CD8⁺ T cell and NK cell reconstitution and also improved graft-versus-leukemia/graft-versus-tumor activity in recipients of haploidentical HSCT. These authors also demonstrated that IL-15 increased intracellular IFN secretion in undivided and slowly proliferating CD8⁺ T cells in recipients of carboxyfluoresceinsuccinimidyl ester–labeled T cells, with no change in the IFN-secreting alloreactive/rapidly proliferative cell population [58].

The efficiency, safety, and feasibility of using IL-15–stimulated CD3/CD19-depleted stem cells or IL-15–activated cytokine-induced killer cells to treat relapse after haploidentical transplantation have been investigated, albeit only in small patient populations [51,57]. Pfeiffer et al. [57] reported that in 8 patients, recovered CD56⁺, CD3⁺/CD4⁺, and CD3⁺/CD8⁺ lymphocyte counts at 15 to 30 days after receipt of IL-15–stimulated stem cell boosts were similar to or even higher than those before the initiation of chemotherapy.

Preemptive Transfer of NK Cells

Adoptive immunotherapy with allogeneic purified NK cell products might exert graft-versus-tumor alloreactivity with little risk of GVHD. In a prospective phase 2 study, Stern et al.

[108] administered purified NK cell products to high-risk patients treated with haploidentical TCD HSCT. Sixteen patients received a total of 29 NK cell infusions on days +3, +40, and +100 post-transplantation. The median dose of infused NK cells per product was $1.21 \times 10^7/\text{kg}$ (range, $0.3\text{--}3.8 \times 10^7/\text{kg}$). Four of the 16 patients were alive at the time of this report, at a median follow-up of 5.8 years. Rapid reconstitution of lymphocyte subsets at day +100 was observed, with a median CD8 T cell count of 266 cells/ μL and a median CD4 T cell count of 181 cells/ μL . These promising results warrant further study to determine the optimal dose and timing of NK cell immunotherapy for antileukemia activity.

Genetically Modified Donor T Cell Infusion

Several researchers have investigated the use of suicide gene–transduced donor T cells to enhance immune reconstitution while controlling GVHD. In a multicenter phase 1/2 clinical trial, purified herpes simplex thymidine kinase suicide gene–transduced donor T cells (TK^{POS} cells) were infused serially into 28 adult patients with a hematologic malignancy after CD34-selected haploidentical transplantation [70]. Twenty-two of the 28 patients who received the purified TK^{POS} cells exhibited a rapid recovery of T cells associated with a concomitant improvement in clinical outcomes and reduced TRM. Patients who did not receive TK^{POS}




Figure 1. A proposed prognostic immune subset-directed strategy to improve immune reconstitution after HIT. On day 0, in vitro alloreactive T cell- and/or B cell-depleted allografts, including NK cells, HSCs, Tregs, and myeloid-derived suppressor cells (MDSCs), were infused to the recipients who received a conditioning regimen to decrease the incidence of GVHD and enhance immune reconstitution. After transplantation, the kinetics of immune recovery were monitored. For patients with rapid immune recovery (as demonstrated by $ALC-30 \geq 300/\mu L$ or another parameter predictive of rapid immune recovery), no interventions were required, and immune reconstitution was monitored only. For patients with delayed immune recovery ($ALC-30 < 300/\mu L$ or other parameters predictive of delayed immune recovery), interventions, including IL-7 and/or adoptive transfer of expanded progenitor T cells or virus-specific donor T cells, were implemented to improve immune reconstitution. MHC, major histocompatibility complex.

cells failed to attain T cell immune reconstitution and experienced dismal outcomes, owing mainly to infectious complications. Impressively, further follow-up of these patients revealed that transfer of TK^{pos} cells not only enhanced the regeneration of $CD3^{+}$ T cells, but also increased counts of the $CD4^{+}$ T cell subset and the T cell subset with a naive phenotype ($CD62L^{+}CD45RA^{+}CD31^{+}$), as well as sjTREC counts, in parallel to the increase in the naive T cell subset. Strikingly, an increase in serum IL-7 level was observed early after the TK^{pos} T cell add-back, followed by a concomitant rise in peripheral T cell counts. Computed tomography-based analysis revealed an increase in thymic tissue even in elderly patients. Additional experiments demonstrated that the antiviral activity of the newly generated T cells was fully maintained.

The biological mechanism responsible for these unexpected findings is unclear. The transient increase in IL-7 might be a contributing factor, as described by Perales et al. [65] in a phase 1 trial, but is likely insufficient to induce long-lasting thymopoiesis. The findings of Vago et al. [70] might suggest new approaches to accelerating immune reconstitution in patients not only after haploidentical transplantation of $CD34^{+}$ stem cells, but also in other transplantation settings, and particularly in elderly patients [2,3,67,71,88].

Adoptive Transfer of Pathogen-Specific T Cells

Gerdemann et al. [49] recently developed a new rapid and simplified manufacturing strategy in which DCs nucleofected with DNA plasmids encoding a range of immunodominant and subdominant viral antigens from adenovirus (Adv), CMV, or Epstein-Barr virus (EBV) are used to generate trivirus-specific T cells. In a recent study, trivirus-directed

T cells were administered to recipients of haploidentical ($n = 5$), matched unrelated ($n = 3$), mismatched unrelated ($n = 1$), or matched related ($n = 1$) transplants with active CMV ($n = 3$), Adv ($n = 1$), EBV ($n = 2$), EBV + Adv ($n = 2$), or CMV + Adv ($n = 2$) infections. The cells produced complete virologic responses in 80% of recipients, including all patients with dual infections. In each case, a decrease in viral load was correlated with an increase in the frequency of T cells directed against the infecting virus(es); both immediate and delayed toxicities were absent [68]. The authors suggested that a DNA plasmid-based approach might increase both the feasibility and applicability of T cell therapy.

Little data are available on vaccine responses in patients undergoing HIT [109]. More recently, several other approaches, including the depletion of naive lymphocytes using Fas ligand [53] and the infusion of $CD62L^{-}$ memory T cells [107], have been used in preclinical models to enhance immune reconstitution after transplantation. Additional studies are warranted to investigate whether these preclinical approaches can be translated into clinical trials.

FUTURE DIRECTIONS

Delayed immune reconstitution is an important issue in haploidentical HSCT. Substantial progress has been made in the field, including insight into recovered immune subsets that can predict outcomes and new strategies for accelerating immune recovery. These advances make it possible to establish a prognostic immune subset-directed strategy to improve immune recovery in haploidentical settings (Figure 1). To achieve this goal, prospective, multicenter trials are needed to investigate whether a low ALC-30 or other simple immune parameters can be used to identify which patients should be treated to improve immune recovery. In

addition, pilot studies should be conducted to evaluate the most effective immune reconstitution-accelerating strategy or combination of approaches. Although promising survival has been achieved with the establishment of many haploidentical transplantation protocols, we believe that a predictive immune subset-guided strategy to accelerate immune recovery might further improve transplantation outcomes in the future, making haploidentical transplantation a routine strategy for patients requiring HSCT.

ACKNOWLEDGMENTS

The authors would like to express their apologies to those whose important work was not cited in this review. They thank San Francisco Edit (www.sefedit.net) for their assistance in editing the manuscript.

Financial disclosure: Support was provided by the Key Program of the National Natural Science Foundation of China (Grant 81230013), the Chang Jiang Scholars Program, and the Beijing Municipal Science and Technology Commission (Grants Z121107002612035 and Z121107002812033).

Conflict of interest statement: There are no conflicts of interest to report.

REFERENCES

- Bashey A, Zhang X, Sizemore CA, et al. T cell-replete HLA-haploidentical hematopoietic transplantation for hematologic malignancies using post-transplantation cyclophosphamide results in outcomes equivalent to those of contemporaneous HLA-matched related and unrelated donor transplantation. *J Clin Oncol*. 2013;31:1310–1316.
- Di Bartolomeo P, Santarone S, De Angelis G, et al. Haploidentical, unmanipulated, G-CSF-primed bone marrow transplantation for patients with high-risk hematologic malignancies. *Blood*. 2013;121:849–857.
- Dvorak CC, Gilman AL, Horn B, et al. Haploidentical related-donor hematopoietic cell transplantation in children using megadoses of CliniMACs-selected CD34(+) cells and a fixed CD3(+) dose. *Bone Marrow Transplant*. 2013;48:508–513.
- Chang Y-J, Huang X-J. Haploidentical bone marrow transplantation without T cell depletion. *Semin Oncol*. 2012;39:653–663.
- Bolanos-Meade J, Fuchs EJ, Luznik L, et al. HLA-haploidentical bone marrow transplantation with posttransplant cyclophosphamide expands the donor pool for patients with sickle cell disease. *Blood*. 2012;120:4285–4291.
- Im HJ, Koh KN, Choi ES, et al. Excellent outcome of haploidentical hematopoietic stem cell transplantation in children and adolescents with acquired severe aplastic anemia. *Biol Blood Marrow Transplant*. 2013;19:754–759.
- Wang Y, Liu DH, Liu KY, et al. Long-term follow-up of haploidentical hematopoietic stem cell transplantation without in vitro T cell depletion for the treatment of leukemia: nine years of experience at a single center. *Cancer*. 2013;119:978–985.
- Federmann B, Bornhauser M, Meisner C, et al. Haploidentical allogeneic hematopoietic cell transplantation in adults using CD3/CD19 depletion and reduced intensity conditioning: a phase II study. *Haematologica*. 2012;97:1523–1531.
- Reisner Y, Hagin D, Martelli MF. Haploidentical hematopoietic transplantation: current status and future perspectives. *Blood*. 2011;118:6006–6017.
- Krenger W, Blazar BR, Hollander GA. Thymic T-cell development in allogeneic stem cell transplantation. *Blood*. 2011;117:6768–6776.
- Geddes M, Storek J. Immune reconstitution following hematopoietic stem-cell transplantation. *Best Pract Res Clin Haematol*. 2007;20:329–348.
- Seggewiss R, Einsele H. Immune reconstitution after allogeneic transplantation and expanding options for immunomodulation: an update. *Blood*. 2010;115:3861–3868.
- Bosch M, Khan FM, Storek J. Immune reconstitution after hematopoietic cell transplantation. *Curr Opin Hematol*. 2012;19:324–335.
- Toubert A, Glauzy S, Douay C, Clave E. Thymus and immune reconstitution after allogeneic hematopoietic stem cell transplantation in humans: never say never again. *Tissue Antigens*. 2012;79:83–89.
- Mackall C, Fry T, Gress R, et al. Background to hematopoietic cell transplantation, including post-transplant immune recovery. *Bone Marrow Transplant*. 2009;44:457–462.
- Pfeiffer MM, Feuchtinger T, Teltschik HM, et al. Reconstitution of natural killer cell receptors influences natural killer activity and relapse rate after haploidentical transplantation of T- and B-cell-depleted grafts in children. *Haematologica*. 2010;95:1381–1388.
- Federmann B, Hagele M, Pfeiffer M, et al. Immune reconstitution after haploidentical hematopoietic cell transplantation: impact of reduced-intensity conditioning and CD3/CD19-depleted grafts. *Leukemia*. 2011;25:121–129.
- Chang Y-J, Zhao X-Y, Huo MR, et al. Influence of lymphocyte recovery on outcome of haploidentical transplantation for hematologic malignancies. *Medicine*. 2009;88:322–330.
- Oevermann L, Lang P, Feuchtinger T, et al. Immune reconstitution and strategies for rebuilding the immune system after haploidentical stem cell transplantation. *Ann NY Acad Sci*. 2012;1266:161–170.
- Isgro A, Marzali M, Sodani P, et al. Immunohematologic reconstitution in pediatric patients after T cell-depleted HLA-haploidentical stem cell transplantation for thalassemia. *Biol Blood Marrow Transplant*. 2010;16:1557–1566.
- Dror Y, Gallagher R, Wara DW, et al. Immune reconstitution in severe combined immunodeficiency disease after lectin-treated, T-cell-depleted haplocompatible bone marrow transplantation. *Blood*. 1993;81:2021–2030.
- Chen X, Hale GA, Barfield R, et al. Rapid immune reconstitution after a reduced-intensity conditioning regimen and a CD3-depleted haploidentical stem cell graft for paediatric refractory haematological malignancies. *Br J Haematol*. 2006;135:524–532.
- Clave E, Lisini D, Douay C, et al. A low thymic function is associated with leukemia relapse in children given T-cell-depleted HLA-haploidentical stem cell transplantation. *Leukemia*. 2012;26:1886–1888.
- Chang Y-J, Zhao X-Y, Huang X-J. Effects of the NK cell recovery on outcomes of unmanipulated haploidentical blood and marrow transplantation for patients with hematologic malignancies. *Biol Blood Marrow Transplant*. 2008;14:323–334.
- Chang Y-J, Zhao X-Y, Huo MR, et al. Immune reconstitution following unmanipulated HLA-mismatched/haploidentical transplantation compared with HLA-identical sibling transplantation. *J Clin Immunol*. 2012;32:268–280.
- Haddad E, Landais P, Friedrich W, et al. Long-term immune reconstitution and outcome after HLA-nonidentical T-cell-depleted bone marrow transplantation for severe combined immunodeficiency: a European retrospective study of 116 patients. *Blood*. 1998;91:3646–3653.
- Friedrich W, Honig M, Schulz A, Muller SM. Immune reconstitution in congenital disorders after HLA-haploidentical hematopoietic stem cell transplantation. *Blood Cells Mol Dis*. 2004;33:291–293.
- Nguyen S, Dhedin N, Vernant JP, et al. NK-cell reconstitution after haploidentical hematopoietic stem-cell transplantations: immaturity of NK cells and inhibitory effect of NKG2A override GvL effect. *Blood*. 2005;105:4135–4142.
- Fu YW, Wu DP, Cen JN, et al. Patterns of T-cell reconstitution by assessment of T-cell receptor excision circle and T-cell receptor clonal repertoire after allogeneic hematopoietic stem cell transplantation in leukemia patients: a study in Chinese patients. *Eur J Haematol*. 2007;79:138–145.
- Rizzieri DA, Koh LP, Long GD, et al. Partially matched, non-myeloablative allogeneic transplantation: clinical outcomes and immune reconstitution. *J Clin Oncol*. 2007;25:690–697.
- Luo XH, Huang X-J, Liu KY, et al. Protective immunity transferred by infusion of cytomegalovirus-specific CD8(+) T cells within donor grafts: its associations with cytomegalovirus reactivation following unmanipulated allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2010;16:994–1004.
- Bader P, Soerensen J, Jarisch A, et al. Rapid immune recovery and low TRM in haploidentical stem cell transplantation in children and adolescence using CD3/CD19-depleted stem cells. *Best Pract Res Clin Haematol*. 2011;24:331–337.
- de Lalla C, Rinaldi A, Montagna D, et al. Invariant NKT cell reconstitution in pediatric leukemia patients given HLA-haploidentical stem cell transplantation defines distinct CD4⁺ and CD4⁺ subset dynamics and correlates with remission state. *J Immunol*. 2011;186:4490–4499.
- Grosso D, Carabasi M, Filicco-O'Hara J, et al. A 2-step approach to myeloablative haploidentical stem cell transplantation: a phase 1/2 trial performed with optimized T-cell dosing. *Blood*. 2011;118:4732–4739.
- Marcanaro E, Carlomagno S, Pesce S, et al. Role of alloreactive KIR2DS1(+) NK cells in haploidentical hematopoietic stem cell transplantation. *J Leukocyte Biol*. 2011;90:661–667.
- Ciurea SO, Mulanovich V, Saliba RM, et al. Improved early outcomes using a T cell-replete graft compared with T cell-depleted haploidentical hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2012;18:1835–1844.
- van Den Brink M, Leen AM, Baird K, et al. Enhancing immune reconstitution: from bench to bedside. *Biol Blood Marrow Transplant*. 2013;19(1 Suppl):S79–S83.
- Aversa F, Terenzi A, Tabilio A, et al. Full haplotype-mismatched hematopoietic stem-cell transplantation: a phase II study in patients

- with acute leukemia at high risk of relapse. *J Clin Oncol*. 2005;23:3447–3454.
39. Ding L, Zheng XL, Dong L, et al. Analysis of T lymphocyte absolute number and function in the early phase after haploidentical hematopoietic stem cell transplantation. *J Exp Hematol*. 2013;21:702–706.
 40. Zhao X-Y, Huang X-J, Liu KY, et al. Prognosis after unmanipulated HLA-haploidentical blood and marrow transplantation is correlated to the numbers of KIR ligands in recipients. *Eur J Haematol*. 2007;78:338–346.
 41. Perez-Martinez A, Gonzalez-Vicent M, Valentín J, et al. Early evaluation of immune reconstitution following allogeneic CD3/CD19-depleted grafts from alternative donors in childhood acute leukemia. *Bone Marrow Transplant*. 2012;47:1419–1427.
 42. Ciurea SO, Mulanovich V, Jiang Y, et al. Lymphocyte recovery predicts outcomes in cord blood and T cell–depleted haploidentical stem cell transplantation. *Biol Blood Marrow Transplant*. 2011;17:1169–1175.
 43. Kesserwan C, Munchel A, Kos F, et al. Immune reconstitution after nonmyeloablative, T-cell–replete, HLA-haploidentical BMT with post-transplantation cyclophosphamide [abstract]. *Biol Blood Marrow Transplant*. 2011;17:S230.
 44. Lang P, Feuchtinger TF, Schlegel P, et al. A new dosing scheme of ATG-F prevents rejection and maintains immune recovery in haploidentical T and B cell–depleted stem cell transplantation. *ASH Annual Meeting Abstracts*. 2012;120:4154.
 45. Azevedo RI, Soares MV, Albuquerque AS, et al. Long-term immune reconstitution of naive and memory T cell pools after haploidentical hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2013;19:703–712.
 46. Jain N, Liu H, Artz AS, et al. Immune reconstitution after combined haploidentical and umbilical cord blood transplant. *Leuk Lymphoma*. 2013;54:1242–1249.
 47. Luo XH, Huang X-J, Li D, et al. Immune reconstitution to cytomegalovirus following partially matched-related donor transplantation: impact of in vivo T-cell depletion and granulocyte colony-stimulating factor–primed peripheral blood/bone marrow mixed grafts. *Transplant Infect Dis*. 2013;15:22–33.
 48. Sanchez-Dominguez R, Pereira-Mendez S, Gomez A, et al. Notch signals contribute to preserve the multipotentiality of human CD34(+)CD38(–)CD45RA(–)CD90(+) hematopoietic progenitors by maintaining T cell lineage differentiation potential. *Exp Hematol*. 2012;40:983–993.e4.
 49. Gerdemann U, Keirnan JM, Katari UL, et al. Rapidly generated multivirus-specific cytotoxic T lymphocytes for the prophylaxis and treatment of viral infections. *Mol Ther*. 2012;20:1622–1632.
 50. Gerdemann U, Vera JF, Rooney CM, Leen AM. Generation of multivirus-specific T cells to prevent/treat viral infections after allogeneic hematopoietic stem cell transplant. *J Vis Exp*. 2011;51:2736.
 51. Rettinger E, Bonig H, Wehner S, et al. Feasibility of IL-15–activated cytokine-induced killer cell infusions after haploidentical stem cell transplantation. *Bone Marrow Transplant*. 2013;48:1141–1143.
 52. Di Ianni M, Falzetti F, Carotti A, et al. Tregs prevent GVHD and promote immune reconstitution in HLA-haploidentical transplantation. *Blood*. 2011;117:3921–3928.
 53. Askenasy N, Mizrahi K, Ash S, et al. Depletion of naive lymphocytes with fas ligand ex vivo prevents graft-versus-host disease without impairing T cell support of engraftment or graft-versus-tumor activity. *Biol Blood Marrow Transplant*. 2013;19:185–195.
 54. Wils EJ, Aerts-Kaya FS, Rombouts EJ, et al. Keratinocyte growth factor and stem cell factor to improve thymopoiesis after autologous CD34⁺ cell transplantation in rhesus macaques. *Biol Blood Marrow Transplant*. 2012;18:55–65.
 55. Innes AJ, Beattie R, Sergeant R, et al. Escalating-dose HLA-mismatched DLI is safe for the treatment of leukaemia relapse following alemtuzumab-based myeloablative allo-SCT. *Bone Marrow Transplant*. 2013;48:1324–1328. <http://dx.doi.org/10.1038/bmt.2013.69>.
 56. Perruccio K, Topini F, Tosti A, et al. Optimizing a photoallodepletion protocol for adoptive immunotherapy after haploidentical SCT. *Bone Marrow Transplant*. 2012;47:1196–1200.
 57. Pfeiffer MM, Schumm M, Muller I, et al. IL-15–stimulated CD3/CD19-depleted stem-cell boosts in relapsed pediatric patients after haploidentical SCT. *Leukemia*. 2012;26:2435–2439.
 58. Sauter CT, Bailey CP, Panis MM, et al. Interleukin-15 administration increases graft-versus-tumor activity in recipients of haploidentical hematopoietic SCT. *Bone Marrow Transplant*. 2013;48:1237–1242.
 59. Chang Y-J, Zhao X-Y, Huo MR, et al. Clinical impact of absolute lymphocyte count on day 30 after unmanipulated haploidentical blood and marrow transplantation for pediatric patients with hematological malignancies. *Am J Hematol*. 2011;86:227–230.
 60. Chang Y-J, Zhao X-Y, Xu LP, et al. Early lymphocyte recovery predicts superior overall survival after unmanipulated haploidentical blood and marrow transplant for myelodysplastic syndrome and acute myeloid leukemia evolving from myelodysplastic syndrome. *Leuk Lymphoma*. 2013;54:2671–2677.
 61. Lilleri D, Gerna G, Zelini P, et al. Monitoring of human cytomegalovirus and virus-specific T-cell response in young patients receiving allogeneic hematopoietic stem cell transplantation. *PLoS ONE*. 2012;7:e41648.
 62. Shimoni A. Haplo-identical stem-cell transplantation: the challenge of immune reconstitution. *Leuk Lymphoma*. 2013;54:2579–2580.
 63. Fujioka T, Tamaki H, Ikegami K, et al. Frequency of CD4(+)FOXP3(+) regulatory T-cells at early stages after HLA-mismatched allogeneic hematopoietic SCT predicts the incidence of acute GVHD. *Bone Marrow Transplant*. 2013;48:859–864.
 64. Feuchtinger T, Opher K, Bethge WA, et al. Adoptive transfer of pp65-specific T cells for the treatment of chemorefractory cytomegalovirus disease or reactivation after haploidentical and matched unrelated stem cell transplantation. *Blood*. 2010;116:4360–4367.
 65. Perales MA, Goldberg JD, Yuan J, et al. Recombinant human interleukin-7 (CYT107) promotes T-cell recovery after allogeneic stem cell transplantation. *Blood*. 2012;120:4882–4891.
 66. Chang Y-J, Huang X-J. Donor lymphocyte infusions for relapse after allogeneic transplantation: when, if and for whom? *Blood Rev*. 2013;27:55–62.
 67. Ciceri F, Bonini C, Stanghellini MT, et al. Infusion of suicide-gene–engineered donor lymphocytes after family haploidentical haemopoietic stem-cell transplantation for leukaemia (the TK007 trial): a non-randomised phase I-II study. *Lancet Oncol*. 2009;10:489–500.
 68. Gerdemann U, Katari UL, Papadopoulos A, et al. Safety and clinical efficacy of rapidly-generated trivirus-directed T cells as treatment for adenovirus, EBV and CMV infections after allogeneic hematopoietic stem cell transplant. *Mol Ther*. 2013;21:2113–2121.
 69. Blyth E, Clancy L, Simms R, et al. Donor-derived CMV-specific T cells reduce the requirement for CMV-directed pharmacotherapy after allogeneic stem cell transplantation. *Blood*. 2013;121:3745–3758.
 70. Vago L, Oliveira G, Bondanza A, et al. T-cell suicide gene therapy prompts thymic renewal in adults after hematopoietic stem cell transplantation. *Blood*. 2012;120:1820–1830.
 71. Luznik L, O'Donnell PV, Symons HJ, et al. HLA-haploidentical bone marrow transplantation for hematologic malignancies using non-myeloablative conditioning and high-dose, posttransplantation cyclophosphamide. *Biol Blood Marrow Transplant*. 2008;14:641–650.
 72. Zhao X-Y, Chang Y-J, Huang X-J. Conflicting impact of alloreactive NK cells on transplantation outcomes after haploidentical transplantation: do the reconstitution kinetics of natural killer cells create these differences? *Biol Blood Marrow Transplant*. 2011;17:1436–1442.
 73. Thoma MD, Huneke TJ, DeCook LJ, et al. Peripheral blood lymphocyte and monocyte recovery and survival in acute leukemia post-myeloablative allogeneic hematopoietic stem cell transplant. *Biol Blood Marrow Transplant*. 2012;18:600–607.
 74. DeCook LJ, Thoma M, Huneke T, et al. Impact of lymphocyte and monocyte recovery on the outcomes of allogeneic hematopoietic SCT with fludarabine and melphalan conditioning. *Bone Marrow Transplant*. 2013;48:708–714.
 75. Chang CC, Wright A, Punnonen J. Monocyte-derived CD11a⁺ and CD11c⁺ dendritic cell subsets differ in their cytokine production profiles, susceptibilities to transfection, and capacities to direct Th cell differentiation. *J Immunol*. 2000;165:3584–3591.
 76. Sun YQ, Xu LP, Liu DH, et al. The incidence and risk factors of invasive fungal infection after haploidentical haematopoietic stem cell transplantation without in vitro T-cell depletion. *Clin Microbiol Infect*. 2012;18:997–1003.
 77. Brennan PJ, Brigl M, Brenner MB. Invariant natural killer T cells: an innate activation scheme linked to diverse effector functions. *Nat Rev Immunol*. 2013;13:101–117.
 78. Chaidos A, Patterson S, Szydlo R, et al. Graft invariant natural killer T-cell dose predicts risk of acute graft-versus-host disease in allogeneic hematopoietic stem cell transplantation. *Blood*. 2012;119:5030–5036.
 79. Clave E, Lisini D, Douay C, et al. Thymic function recovery after unrelated donor cord blood or T-cell–depleted HLA-haploidentical stem cell transplantation correlates with leukemia relapse. *Front Immunol*. 2013;4:54.
 80. Sivori S, Carlomagno S, Falco M, et al. Natural killer cells expressing the KIR2DS1-activating receptor efficiently kill T-cell blasts and dendritic cells: implications in haploidentical HSCT. *Blood*. 2011;117:4284–4292.
 81. Xiao TZ, Singh K, Dunn E, et al. T cell and B cell immunity can be reconstituted with mismatched hematopoietic stem cell transplantation without alkylator therapy in artemis-deficient mice using anti-natural killer cell antibody and photochemically treated sensitized donor T cells. *Biol Blood Marrow Transplant*. 2012;18:200–209.
 82. Ruggeri L, Capanni M, Urbani E, et al. Effectiveness of donor natural killer cell alloreactivity in mismatched hematopoietic transplants. *Science*. 2002;295:2097–2100.
 83. Handgretinger R. Negative depletion of CD3⁺ and TCRαβ⁺ T cells. *Curr Opin Hematol*. 2012;19:434–439.
 84. Xie M, Fu HX, Chang Y-J, et al. Characteristics and influencing factors of CD19⁺ B cell reconstitution in patients following haploidentical/

- mismatched hematopoietic stem cell transplantation. *Int J Hematol*. 2012;96:109–121.
85. Huang X-J, Xu L-P, Liu K-Y, et al. Partially matched related donor transplantation can achieve outcomes comparable with unrelated donor transplantation for patients with hematologic malignancies. *Clin Cancer Res*. 2009;15:4777–4783.
 86. Huang X-J, Liu DH, Liu K-Y, et al. Treatment of acute leukemia with unmanipulated HLA-mismatched/haploidentical blood and bone marrow transplantation. *Biol Blood Marrow Transplant*. 2009;15:257–265.
 87. Lu DP, Dong L, Wu T, et al. Conditioning including antithymocyte globulin followed by unmanipulated HLA-mismatched/haploidentical blood and marrow transplantation can achieve comparable outcomes with HLA-identical sibling transplantation. *Blood*. 2006;107:3065–3073.
 88. Raiola AM, Dominietto A, Ghiso A, et al. Unmanipulated haploidentical bone marrow transplantation and posttransplantation cyclophosphamide for hematologic malignancies after myeloablative conditioning. *Biol Blood Marrow Transplant*. 2013;19:117–122.
 89. Marin IGG, Perez-Corral AM, Gayoso J, et al. Natural killer (NK) cell reconstitution after haploidentical unmanipulated bone marrow transplantation with reduced-intensity conditioning. ASH Annual Meeting Abstracts. *Blood*. 2011;118:4557.
 90. Koehl U, Bochennek K, Zimmermann SY, et al. Immune recovery in children undergoing allogeneic stem cell transplantation: absolute CD8⁺ CD3⁺ count reconstitution is associated with survival. *Bone Marrow Transplant*. 2007;39:269–278.
 91. Noviello M, Forcina A, Stanghellini MTL, et al. Early reconstitution of T-cell immunity to CMV after HLA-haploidentical hematopoietic stem cell transplantation is a strong surrogate biomarker for lower non-relapse mortality rates. ASH Annual Meeting Abstracts. *Blood*. 2012;120:4191.
 92. Chang Y-J, Huang X-J. Haploidentical hematopoietic stem cell transplantation with unmanipulated granulocyte colony-stimulating factor—mobilized marrow and blood grafts. *Curr Opin Hematol*. 2012;19:454–461.
 93. Dominietto A, Raiola AM, Bruno B, et al. Rapid immune reconstitution following unmanipulated haploidentical BMT with post-transplant high-dose cyclophosphamide ASH Annual Meeting Abstracts. *Blood*. 2011;118:3050.
 94. Merindol N, Champagne MA, Duval M, Soudeyins H. CD8(+) T-cell reconstitution in recipients of umbilical cord blood transplantation and characteristics associated with leukemic relapse. *Blood*. 2011;118:4480–4488.
 95. McIver ZA, Melenhorst JJ, Grim A, et al. Immune reconstitution in recipients of photodepleted HLA-identical sibling donor stem cell transplantations: T cell subset frequencies predict outcome. *Biol Blood Marrow Transplant*. 2011;17:1846–1854.
 96. Gratama JW, Boeckh M, Nakamura R, et al. Immune monitoring with iTag MHC tetramers for prediction of recurrent or persistent cytomegalovirus infection or disease in allogeneic hematopoietic stem cell transplant recipients: a prospective multicenter study. *Blood*. 2010;116:1655–1662.
 97. Wang Y, Liu DH, Xu LP, et al. Superior graft-versus-leukemia effect associated with transplantation of haploidentical compared with HLA-identical sibling donor grafts for high-risk acute leukemia: an historic comparison. *Biol Blood Marrow Transplant*. 2011;17:821–830.
 98. Yan CH, Liu DH, Liu KY, et al. Risk stratification—directed donor lymphocyte infusion could reduce relapse of standard-risk acute leukemia patients after allogeneic hematopoietic stem cell transplantation. *Blood*. 2012;119:3256–3262.
 99. Zhao XS, Jin S, Zhu HH, et al. Wilms' tumor gene 1 expression: an independent acute leukemia prognostic indicator following allogeneic hematopoietic SCT. *Bone Marrow Transplant*. 2012;47:499–507.
 100. Zhao XS, Liu YR, Zhu HH, et al. Monitoring MRD with flow cytometry: an effective method to predict relapse for ALL patients after allogeneic hematopoietic stem cell transplantation. *Ann Hematol*. 2012;91:183–192.
 101. Szabolcs P, Niedzwiecki D. Immune reconstitution after unrelated cord blood transplantation. *Cytotherapy*. 2007;9:111–122.
 102. Brown JA, Stevenson K, Kim HT, et al. Clearance of CMV viremia and survival after double umbilical cord blood transplantation in adults depends on reconstitution of thymopoiesis. *Blood*. 2010;115:4111–4119.
 103. Sauter C, Abboud M, Jia X, et al. Serious infection risk and immune recovery after double-unit cord blood transplantation without antithymocyte globulin. *Biol Blood Marrow Transplant*. 2011;17:1460–1471.
 104. June CH, Blazar BR, Riley JL. Engineering lymphocyte subsets: tools, trials and tribulations. *Nat Rev Immunol*. 2009;9:704–716.
 105. Oliveira G, Greco R, Lupo-Stanghellini MT, et al. Use of TK-cells in haploidentical hematopoietic stem cell transplantation. *Curr Opin Hematol*. 2012;19:427–433.
 106. Li HW, Sykes M. Emerging concepts in haematopoietic cell transplantation. *Nat Rev Immunol*. 2012;12:403–416.
 107. Zhang J, Barefoot BE, Mo W, et al. CD62L⁺ memory T cells enhance T-cell regeneration after allogeneic stem cell transplantation by eliminating host resistance in mice. *Blood*. 2012;119:6344–6353.
 108. Stern M, Passweg JR, Meyer-Monard S, et al. Pre-emptive immunotherapy with purified natural killer cells after haploidentical SCT: a prospective phase II study in two centers. *Bone Marrow Transplant*. 2013;48:433–438.
 109. Chou JF, Kernan NA, Prockop S, et al. Safety and immunogenicity of the live attenuated varicella vaccine following T cell—replete or T cell-depleted related and unrelated allogeneic hematopoietic cell transplantation (alloHCT). *Biol Blood Marrow Transplant*. 2011;17:1708–1713.